



King's Research Portal

DOI:

[10.1042/BST20170508](https://doi.org/10.1042/BST20170508)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Aksentijevic, D., O'Brien, B. A., Eykyn, T. R., & Shattock, M. J. (2018). Is there a causal link between intracellular na elevation and metabolic remodelling in cardiac hypertrophy? *Biochemical Society Transactions*, 46(4), 817-827. <https://doi.org/10.1042/BST20170508>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Review Article

Is there a causal link between intracellular Na_i elevation and metabolic remodelling in cardiac hypertrophy?

Dunja Aksentijevic^{1,2}, Brett A. O'Brien^{2,3}, Thomas R. Eykyn³ and Michael J. Shattock²

¹School of Biological and Chemical Sciences, Queen Mary University of London, G.E. Fogg Building, London, U.K.; ²King's College London, School of Cardiovascular and Medical Sciences, British Heart Foundation Centre of Research Excellence, St Thomas Hospital, London, U.K.; ³Department of Imaging Chemistry and Biology, School of Biomedical Engineering and Imaging Sciences, King's College London, St Thomas' Hospital, London, U.K.

Correspondence: Dunja Aksentijevic (d.aksentijevic@qmul.ac.uk)



Alterations in excitation–contraction coupling and elevated intracellular sodium (Na_i) are hallmarks of pathological cardiac remodelling that underline contractile dysfunction. In addition, changes in cardiac metabolism are observed in cardiac hypertrophy and heart failure (HF) that lead to a mismatch in ATP supply and demand, contributing to poor prognosis. A link between Na_i and altered metabolism has been proposed but is not well understood. Many mitochondrial enzymes are stimulated by mitochondrial calcium (Ca_{mito}) during contraction, thereby sustaining production of reducing equivalents to maintain ATP supply. This stimulation is thought to be perturbed when cytosolic Na_i is high due to increased Ca_{mito} efflux, potentially compromising ATP_{mito} production and leading to metabolic dysregulation. Increased Na_i has been previously shown to affect Ca_{mito}; however, whether Na_i elevation plays a causative role in energetic mismatching in the hypertrophied and failing heart remains unknown. In this review, we discuss the relationship between elevated Na_i, NaK ATPase dysregulation and the metabolic phenotype in the contexts of pathological hypertrophy and HF and their link to metabolic flexibility, capacity (reserve) and efficiency that are governed by intracellular ion homeostasis. The development of non-invasive analytical techniques using nuclear magnetic resonance able to probe metabolism *in situ* in the functioning heart will enable a better understanding of the underlying mechanisms of Na_i overload in cardiac pathophysiology. They will lead to novel insights that help to explain the metabolic contribution towards these diseases, the incomplete rescue observed with current therapies and a rationale for future energy-targeted therapies.

Introduction

Cardiovascular disease is the leading cause of mortality worldwide with its incidence projected to rise significantly in the immediate future. There is a clear need for improved understanding of underlying cellular mechanisms which can aid the development of more effective treatments as well as novel techniques for early diagnosis. There is a convincing evidence that myocardial intracellular Na (Na_i) overload along with metabolic derangement are two important and interconnected pathophysiological features of hypertrophy and heart failure (HF). Na ion homeostasis is regulated by many transporters and membrane pumps [1]. Na/K ATPase (NKA) and its key regulatory protein phospholemman (PLM) play a crucial role in cardiomyocyte transmembrane ion transport and contractility, such that transgenic PLM^{3SA} mice, in which PLM is rendered unphosphorylatable, have chronically elevated Na_i and an increased susceptibility to hypertrophy-induced dysfunction [2]. In addition, due to the high ATP demand required for pump activity and its sarcolemmal localization, there is evidence for an association between NKA and metabolism with its ATP supply thought to be supplied preferentially

Received: 11 April 2018
Revised: 23 May 2018
Accepted: 24 May 2018

Version of Record published:
3 July 2018

by glycolysis [3–5]. More recent studies have suggested that cytosolic Na regulation also plays an important role in linking mitochondrial Ca-dependent ATP production to mechanical activity and ATP demand due to contractile work [6,8]. Nevertheless, the extent that these metabolic alterations (mismatch in ATP supply–demand) reflect chronic cellular remodelling or arise as a consequence of Na_i elevation is not well understood.

Na pump and Na_i regulation in cardiac hypertrophy

In most larger mammalian hearts, with a long action potential, Na_i is maintained at ~4–8 mM [9,10]. In murinae (rats and mice), intracellular Na is significantly elevated (10–20 mM) and this elevated Na is associated with many other adaptations in excitation–contraction (EC) coupling, including a short action potential, a larger recirculating Ca fraction, a dependence on SR Ca release, reduced NCX (sarcolemmal sodium–calcium exchanger) activity, rest potentiation and a negative force–frequency staircase [6].

The cell exploits the energy in the transmembrane Na gradient to drive a plethora of Na-dependent membrane transporters moving ions, substrates, amino acids etc. either into (co-transporters/symports) or out of (exchangers/antiports) the cell (Figure 1). The importance of this trans-sarcolemmal inward Na gradient means that its dissipation in various pathologies such as ischaemia/reperfusion [11], hypertrophy or HF [12,13] is highly detrimental. While some of the Na transport processes are electro-neutral, some are electrogenic and hence both respond to, and contribute to, the membrane potential. Most notably, voltage-gated Na channels are crucially important in generating the upstroke of the cardiac action potential. While there are a large number of Na influx pathways, there is only a single quantitatively significant Na efflux pathway responsible for maintaining the transmembrane Na gradient — the Na/K ATPase or Na/K pump (NKA) [14].

The activity of the NKA is regulated by FXD1, or PLM, the principal sarcolemmal target of protein kinases A and C (Figure 2) [15]. As such, PLM is required for the dynamic control of Na_i during increases in heart rate or during disease and plays a vital role in Na regulation during ‘fight or flight’ [14]. Under physiological conditions, NKA is the only quantitatively significant efflux pathway of Na out of the myocyte (NCX and Na/HCO₃/Cl symporter, in principle, can reverse and efflux Na) [16] (Figure 1).

A hallmark of cardiac hypertrophy and failure is an elevation of Na_i. There is an abundant literature on this phenomenon, although absolute values of measured Na_i are often dissimilar, probably owing to methodological differences as summarized in Table 1. Elevation in Na_i may contribute to the negative force–frequency relationship, slowed relaxation and arrhythmias [17]. While a component of the elevation of Na_i may reflect an increase in Na influx [10], there is a large body of evidence showing that Na/K pump function may also be compromised [2,12,13,18]. Specifically, in cardiac hypertrophy, many studies have shown that NKA pump function, and/or expression, is reduced [2,12,13,18,19]. Cardiac Na_i can also be elevated by several other factors such as hypothermia [16] or by increased intracellular pH via enhanced NHE (sodium–proton exchanger) activity, for example, following ischaemia–reperfusion injury [20].

ATP supply–demand matching in the heart

Fine control of ATP-generating pathways in mitochondria and cytosol are critical to meet the energy demands of cardiac muscle. Supply must be matched to demand as failure to provide an adequate amount of ATP causes a decrease in cellular free energy leading to mechanical failure. The heart utilizes more energy than any other organ — with 2% of its total ATP reserves consumed per beat, it turns over its total ATP pool in less than 1 min and utilizes 6 kg of ATP every day [21–23]. This enormous energy demand is related primarily to ATP-dependent processes driving EC coupling [24]. About 70–75% of total intracellular ATP is used for force generation powering work output, with the remaining 25–30% is used for basal metabolism [25–27]. In terms of force generation, it is estimated that the actomyosin ATPase accounts for 76%, SERCA (sarcoendoplasmic reticulum Ca²⁺ ATPase) 15% and NKA for 9% of ATP utilization [27].

To synthesize the ATP required for normal function, the adult heart converts chemical energy primarily stored in free fatty acids (FFAs) (60–90%) and pyruvate (derived from glucose and lactate 10–40%) into mechanical energy for contraction [28]. The delivery of metabolic substrates, their selection, uptake and oxidation to generate acetyl-CoA for tricarboxylic acid (TCA) cycle entry and ATP generation in the electron transport chain (ETC) comprises three stages of myocardial ATP supply as summarized in Figure 1. However, cardiac workload varies constantly, including several-fold increase in cardiac output during exercise, thus requiring rapid and continuous matching of ATP supply to demand. This renders the heart a metabolic omnivore, giving it a high degree of substrate flexibility to rapidly switch substrate preference and utilization [28]. The apparent




Figure 1. Major Na⁺ influx and efflux pathways and metabolic pathways involved in ATP supply.

The delivery of metabolic substrates, their selection and uptake are followed by OXPHOS. It involves electron shuttling from cytosolic to mitochondrial reducing equivalents, transfer of energy by electrons from reducing equivalents to ETC complexes and generation of electrochemical proton (H^+) gradient within the mitochondrial intermembrane space (respiratory complexes I, II, III, IV). The release of H^+ gradient is coupled to the synthesis of ATP from ADP + P_i by F_0F_1 -ATPase (complex V), contributing >95% of ATP synthesis under aerobic conditions. The final stage of myocardial ATP supply (phosphotransfer) involves delivery of ATP from mitochondria to sites of use. This involves ADP-ATP exchange across the inner mitochondrial membrane by the adenine nucleotide transporter (ANT) and propagation of local ATP/ADP disequilibria primarily by the creatine kinase (CK). Abbreviations: TAG, triacylglycerol; PCr, phosphocreatine; ANT, adenine nucleotide transporter; GLUT, glucose transporter; CD36, fatty acid transporter; PPP, pentose phosphate pathway; LDH, lactate dehydrogenase; PDH, pyruvate dehydrogenase; CPT, carnitine palmitoyltransferase; CACT, carnitine-acylcarnitine translocase; MCU, mitochondrial calcium uniporter; α -KDH, α -ketoglutarate dehydrogenase; IDH, isocitrate dehydrogenase; mitoCK, mitochondrial creatine kinase; IMM, inner mitochondrial membrane; OMM, outer mitochondrial membrane; Q, quinone pool; c, cytochrome c; MPC, mitochondrial pyruvate carrier; e^- , electrons; CGP, mitochondrial Na-Ca exchanger inhibitor CGP-37157. *Mitochondrial calcium-sensitive dehydrogenases (pyruvate dehydrogenase, isocitrate dehydrogenase and α -ketoglutarate dehydrogenase).

opposing relationship between carbohydrates and FFAs in the heart is, in part, due to the Randle (glucose–fatty acid) cycle, thus optimizing energy supply by avoiding energetic inefficiency and ‘waste’ [29].




Figure 2. Schematic depiction of a structure–function relationship (regulation) between PLM and Na pump.

The cytoplasmic tail of unphosphorylated PLM interacts closely with the membrane and α -subunit of Na pump, whereas phosphorylation alters the association between the pump and PLM by moving the cytosolic arm away from the pump, but not by promoting their dissociation. Phosphorylation or ablation of PLM relieves inhibition of the Na pump by increasing its V_{\max} and apparent Na affinity. Under stress, phosphorylation of PLM allows the heart to reduce its Na and Ca load and prevents lethal arrhythmias. Adapted from Pavlovic et al. [15].

The failing heart

First identified in the early 20th century, and now a well-established energy starvation hypothesis, it is proposed that maladaptive metabolic remodelling precedes, initiates and maintains adverse contractile dysfunction in hypertrophy and HF [23,24]. Advances in analytical technologies and understanding of metabolic mechanisms have improved our insights into the phenomenon and helped to classify metabolic alterations leading to myocardial energy starvation into those related to substrate utilization, intermediary metabolism and energetics. Using *in vivo* ^{31}P nuclear magnetic resonance (NMR), Neubauer [23] found that the myocardial phosphocreatine-to-ATP ratio (PCr:ATP) can be used as a reliable prognostic indicator of dilated cardiomyopathy (DCM) where 44% of DCM patients with a PCr:ATP of <1.6 died of cardiovascular causes vs. 5% with a PCr:ATP of >1.6. Cardiac hypertrophy induces a switch in substrate utilization from dominant FFA oxidation towards carbohydrate utilization which is similar to the foetal metabolic phenotype [24,30–32]. The onset of this switch (and thereby the stage at which it could potentially be targeted therapeutically) is currently debated as numerous studies suggest that ATP levels are sustained during the early stages of remodelling and only decrease (30–40%) during advanced stages of HF [33–37]. There have also been numerous preclinical studies as well as clinical data inferring mitochondrial respiratory impairment (complex activities and/or altered expression of the ETC complexes, ATP synthase and adenine nucleotide translocase) in hypertrophy and HF [38–40].

EC coupling, specifically Ca handling by SERCA, has also been linked to the time course of metabolic alterations during hypertrophy: SERCA preferentially uses glycolytically derived ATP over OXPHOS (oxidative phosphorylation) [41] and therefore switching to a more glycolytic phenotype during hypertrophy, and HF could reflect increased SERCA activity to sustain adequate Ca homeostasis. NKA pump also requires glycolysis for normal Na homeostasis, potentially due to preferential fuelling of NKA by cytosolic glycolytically derived ATP and its spatial proximity to the pump [4,42,43]. However, the substrate switch and energetic deficit alone cannot explain Na accumulation observed in hypertrophy and failure. The debate is similar to the arguments about Na elevation in ischaemia and revolves around energetic inhibition of the NKA: the substrate switch from fatty acids to glucose leads to impaired energetic reserve and decline in cytosolic ATP, thus limiting the energy supply to the pump leading to Na accumulation. However, it has been previously shown that even during severe metabolic stress such as ischaemia, intracellular Na rises at a time when the total ATP

Table 1 Summary of studies quantifying bulk cytosolic $[Na]_i$ in the myocyte under both physiological and pathophysiological conditions across various mammalian species

Species	$[Na]_i$ (mM)	Method used	Reference
Human	8.0	SBFI (sodium-binding benzofuran isophthalate)-loaded muscle strips paced at 0.25 Hz	[17]
Human LVH	14.2	Na-selective microelectrodes; muscle strips at rest	[67]
Human failing	12.1	SBFI-loaded muscle strips paced at 0.25 Hz	[17]
Human MVD	11.8	Na-selective microelectrodes; muscle strips at rest	[67]
Sheep	5–6.4 5.8–7.9	Na-selective microelectrodes; Purkinje fibres at 1 Hz and at rest Na-selective microelectrodes; muscle strips at rest	[68,69] [70]
Dog	8.9–10.4	Na-selective microelectrodes; Purkinje fibres at rest and 1 Hz	[71]
Guinea pig	4.7–8.0 6.4 5.1–5.2	Na-selective microelectrodes; muscle strips at rest ^{23}Na NMR; isolated perfused heart SBFI-loaded myocytes at rest	[67,70,72,73,74] [75] [9,76]
Guinea pig	12.1	Na-selective microelectrodes; muscle strips at rest	[67]
LVH	12.8	^{23}Na NMR; isolated perfused heart	[75]
Guinea pig failing	16.8	SBFI-loaded myocytes at rest	[77]
Ferret	7.8	Na-selective microelectrodes; muscle strips at rest	[78]
Ferret RVH	8.0	Na-selective microelectrodes; muscle strips at rest	[78]
Rabbit	7.2 3.8–4.5	Na-selective microelectrodes; muscle strips at 0.5 Hz SBFI-loaded myocytes at rest	[6,10] [10,79]
Rat	12.7 8.5–30 5.1–21 17.5	Na-selective microelectrodes; muscle strips at 0.5 Hz Na-selective microelectrodes; myocytes at rest SBFI-loaded myocytes at rest ^{23}Na NMR; isolated perfused arrested hearts	[6] [80] [9,10,79,81] [82]
Mouse	11.6	^{23}Na NMR; isolated perfused heart	[51]
Mouse	14	SBFI-loaded myocytes at rest	[2]
Mouse LVH	23	SBFI-loaded myocytes at rest	[2]

concentration greatly exceeds the K_m for the pump (~ 0.1 – 0.8 mmol/l) and the free energy of ATP exceeds that required for pump activity (~ 44 kJ/mol) [44].

Myocardial Na_i elevation and metabolic remodelling: the chicken or the egg?

In spite of significant evidence to support the concomitance of Na_i overload and metabolic remodelling during cardiac hypertrophy and HF, there have been very few studies investigating the interaction between these pathophysiological events. Using isolated rat mitochondria, Iwai et al. [45] demonstrated that increasing extramitochondrial Na (Na_{ex}) from physiological (12.5 mM) to supraphysiological (≥ 25 mM) concentrations significantly reduced state 3 respiration, suggesting reduced mitochondrial ATP supply as well as reduced mitochondrial membrane potential. However, the present study offered no insights into the mechanism underlying the effect of Na_i overload on whole cell metabolism.

A series of studies focusing on the mitochondrial transport of Na and Ca and its relationship with mitochondrial ATP production showed a stimulation of mitochondrial ATP production by Ca; however, the mitochondrial Ca transport kinetics and its regulation by Na_i are still not completely understood [45–48]. The majority of Ca_{mito} uptake is by the Ca_{mito} uniporter (MCU), while the Na/ Ca_{mito} exchanger (NCLX) is thought to be the predominant mechanism for Ca extrusion [49] (Figure 1). The impact of Na_i on Ca_m has been elucidated by Cox and Matlib [46] using fura-2 to measure Ca_m in isolated cardiac mitochondria from healthy rabbits. Mitochondria incubated with increasing concentrations of Na_{ex} using NaCl in the physiological range

showed reduced Ca_{mito} as well as reduced NADH production and state 3 respiration. On the other hand, inhibition of NCLX with three inhibitors (from highest to lowest potency: CGP-37157 > clonazepam > D-cis-diltiazem) and the MCU inhibitor ruthenium red substantially increased Ca_{m} , NADH production and state 3 respiration in a dose-dependent manner. This study supported the findings of Iwai et al. [45] and the hypothesis that Na_i overload dysregulates ATP supply–demand matching.

However, this study did not provide information on the beat-to-beat kinetics of Ca_{m} transport and its relation to mitochondrial energy production.

Isolated mitochondria experiments should be treated with caution, given the measurements are performed in the absence of important ATP sinks (myosin ATPase, NKA and SERCA) and substrate utilization pathways (glycolysis and β -oxidation). More recently, Maack et al. [50] used isolated guinea pig cardiomyocytes to measure Ca_i and Ca_{m} during systole and diastole as well as NADH, thereby providing insights into the beat-to-beat regulation of Ca_{m} during increased Na_i . This study showed that both systolic and diastolic Ca_{m} are significantly reduced by Na_i elevation. Correspondingly, the percentage of NAD(H) in the reduced form was maintained at ~62% in the control group, but was significantly lower in the high Na_i group. In spite of these changes in Ca and [NADH], Na_i elevation did not affect the mitochondrial membrane potential ($\Delta\Psi_{\text{m}}$). Furthermore, NCLX inhibition by CGP-37157 was shown to significantly elevate diastolic Ca_{m} . As these effects were not altered by ruthenium red inhibition of the MCU, it is likely a consequence of increased Ca_{m} extrusion via NCLX on a beat-to-beat basis. The outcome of this study further supported the argument that Na_i is an important regulator of cardiac bioenergetics. However, it remains unclear whether this is truly reflective of a regulatory mechanism in the beating heart and, if so, which metabolic pathways are most affected by Na_i overload.

Measuring Na_i overload in the beating heart

The studies examining the impact of Na_i on mitochondrial ATP provision published to date are subject to major experimental caveats, thus making direct mechanistic translation to *in situ* perfused and *in vivo* myocardium difficult. Specifically, these studies lack integrated experimental approaches as they have been limited to isolated organelles and cells at subphysiological temperatures with limited metabolic readouts.

To elucidate the importance of the link between Na_i and ATP supply–demand matching in the beating heart, experimental models are required either *ex vivo* or *in vivo* where the heart is perfused under physiologically relevant conditions and where Na_i elevation can be induced and reliably measured. To elucidate concomitant changes in substrate metabolism or energetics, it is also necessary to be able to quantify a wide range of metabolites involved in energy homeostasis in these models. We have previously applied and validated techniques able to measure intracellular Na_i in the Langendorff perfused mouse [51,52] or rat heart preparations [53] using NMR.

^{23}Na NMR has historically been used to distinguish the small intra Na_i versus large extracellular Na_e pools employing paramagnetic shift reagents such as Tm(DOTP) [thulium (III) 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetra(methylenephosphonate)] [51] to separate the two. However, these reagents are efficient chelators of Ca and Mg leading to modified ion homeostasis and reduced cardiac contractility [51]. As a result, shift reagents exhibit significant toxicity precluding their use *in vivo* and question their validity for measuring Na_i *ex vivo*. In contrast, multiple quantum-filtered ^{23}Na NMR, which exploits the quadrupolar property of the ^{23}Na nucleus, has shown potential to probe intra and extracellular pools of Na in the *absence* of shift reagent and therefore under more physiological conditions [54,55]. We have previously investigated the use of these techniques in the perfused mouse heart where we were able to measure elevated intracellular Na in response to the cardiac glycoside ouabain as well as in response to modified buffer compositions, for example, in the absence of K, Ca or Mg. We were further able to verify previous studies showing that the PLM^{3SA} mouse has a chronic elevation of basal Na_i compared with wild-type hearts. Crucially, NMR is also able to probe cardiac energetics by ^{31}P NMR in the same hearts where it is possible to measure the concentrations of ATP, PCr, P_i , intracellular pH as well as PCr:ATP ratio and thereby derive estimates of the Gibbs free energy. Newly emerging techniques such as metabolomics also enable end-point measurements of metabolites in snap frozen extracted myocardial tissue and coronary effluent using either high-resolution NMR or mass spectrometry [53,56,57].

Figure 3 shows the example NMR spectra of two different Langendorff perfused mouse hearts acquired using our previously reported NMR protocols. The left-hand panel displays data from a wild-type control mouse heart with normal baseline Na_i , while the right-hand panel displays data from a hypertrophic mouse heart subject to aortic constriction [2]. The dry weight of the control heart was 30 mg while that of the banded




Figure 3. Representative ^{31}P NMR spectra, triple-quantum-filtered ^{23}Na and conventional 1D ^{23}Na NMR spectra from perfused control and hypertrophied mouse hearts.

The spectra displayed in the left panel (**a**, **c** and **e**) are from a control heart, while those displayed in the right panel (**b**, **d** and **f**) are from a hypertrophied heart. All NMR data were acquired as previously described [51] using a Bruker Avance III 400 MHz wide-bore spectrometer. Briefly, **a** and **b** show ^{31}P spectra, **c** and **d** show triple-quantum-filtered (TQF) ^{23}Na NMR spectra, while **e** and **f** show conventional single-quantum ^{23}Na NMR spectra acquired at the end of the perfusion during infusion of 5 mM Tm(DOTP).

heart was 58 mg measured at the end of the experiment. **Figure 3a,b** shows the ^{31}P NMR spectra acquired with baseline function. **Figure 3c,d** shows the triple-quantum-filtered ^{23}Na NMR spectra acquired with baseline function. **Figure 3e,f** shows the conventional single-quantum-filtered ^{23}Na NMR spectra acquired at the end of the experiment following infusion of 5 mM Tm(DOTP) to shift the large extracellular Na_e signal and enable quantification of the small intracellular Na_i signal. Our previous work suggested that the TQF signal in **Figure 3c,d** consists of a contribution both from the intracellular and extracellular pools of Na, but that the large bulk isotropic signal from the buffer is largely suppressed. These experiments highlight the ability of such NMR techniques to probe both cardiac energetics using ^{31}P NMR and Na_i using triple-quantum-filtered ^{23}Na NMR in the same preparation [51]. The data presented here also highlight experimental challenges in quantifying Na_i in these hearts. Total myocardial Na_i is clearly elevated under conditions of hypertrophy; however, so too is the tissue mass and intracellular volume [51]. Absolute quantification of such data is subject to many experimental assumptions including a phenomenological scaling factor for the NMR observability of Na_i and a scaling factor to estimate intracellular volume. Despite obvious limitations in the methodology, NMR offers unique insights into Na ion homeostasis and cardiac energetics under both physiological and pathophysiological conditions. Additionally, there has resurgence in interest applying MRI techniques for imaging Na distribution *in vivo* [58]. ^{23}Na is the second most sensitive nucleus for *in vivo* detection by NMR after ^1H ; however, sensitivity and spatial resolution remain an issue as well as the ability to separate intra- versus extracellular pools of Na which is also challenging.

Therapeutic potential

Na_i has inadvertently been a known therapeutic target in HF for the last 200 years, and the established example of *in vivo* use of Na_i modulation is the administration of cardiac glycosides (such as digoxin) which are potent inhibitors of NKA. Cardiac glycosides elevate Na_i and lead to a positive inotropic response (due to

release of Ca) that can vary considerably between species. They are steroidal-like compounds found endogenously under normal conditions (e.g. ouabain, digoxin and bufalin) and are elevated in patients with renal failure [59] and HF [60]. However, their clinical use for the treatment of HF is a cautionary tale and limited due to their energetically costly as well as pro-arrhythmic properties [61]. Nevertheless, they remain useful tool for elevating Na_i in experimental models [51–53,56,62]. In spite of the substantial *in vitro* and preclinical evidence to support the targeting of the substrate switch therapeutically, there has been limited successes that have been translated into the clinic. For example, sodium dichloroacetate (pyruvate dehydrogenase kinase inhibitor) appeared to improve contractile performance in 10 HF patients, but a vehicle control group was not included in this study [63,64]. Trimetazadine is currently prescribed for longer term inhibition of FFA oxidation and has been shown to reduce angina and improve cardiac function in patients with DCM [65,66], although these improvements were modest. Given the limited clinical success of targeting substrate utilization to date, it is important to continue to evaluate the potential of targeting other aspects of cardiac metabolism, such as intermediary pathways leading to ATP supply. This could also help identify the role metabolic remodelling plays in transition from pathological hypertrophy towards HF. The question remains whether early prevention of myocardial Na_i elevation could either prevent the origin or alter the course of metabolic derangement in pathological hypertrophy leading to energy starvation and cardiac death. This hypothesis warrants further study including the ongoing development of therapeutics that target these interconnected pathophysiological events.

Abbreviations

Ca_{mito} , mitochondrial calcium; DCM, dilated cardiomyopathy; EC, excitation–contraction; ETC, electron transport chain; FFA, free fatty acids; HF, heart failure; MCU, mitochondrial calcium uniporter; Na_{ex} , extramitochondrial sodium; Na_i , intracellular sodium; NCLX, mitochondrial sodium–calcium exchanger; NCX, sarcolemmal sodium–calcium exchanger; NKA, sodium/potassium ATPase; NMR, nuclear magnetic resonance; OXPHOS, oxidative phosphorylation; PCr:ATP, phosphocreatine-to-ATP ratio; PLM, phospholemman; SBFI, sodium-binding benzofuran isophthalate; SERCA, sarcoendoplasmic reticulum Ca^{2+} ATPase; Tm(DOTP), thulium (III) 1,4,7,10-Tetraazacyclododecane-1,4,7,10-tetra(methylenephosphonate).

Author Contribution

D.A., T.R.E., B.A.O.B. and M.J.S. conceived and designed research. D.A. drafted the manuscript. D.A. and T.R.E. prepared figures. D.A., B.A.O.B., T.R.E. and M.J.S. edited, revised the manuscript and approved the final version of the manuscript.

Funding

This work was supported by a British Heart Foundation Programme Grant [RG/12/4/29426] to M.J.S. and King's College London British Heart Foundation Centre of Research Excellence [RE/08/003]. The research was supported by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London; the Centre of Excellence in Medical Engineering funded by the Wellcome Trust and EPSRC under grant number [WT 088641/Z/09/Z] and the King's College London and UCL Comprehensive Cancer Imaging Centre funded by the CRUK and EPSRC in association with the MRC and DoH (England). The views expressed are those of the author and not necessarily those of the NHS, the NIHR or the Department of Health.

Acknowledgements

The authors thank Anja Karlstaedt, Heinrich Taegtmeyer, David Sanchez-Tatay, Marina Basalay, Andrew Atkinson, Alpesh Thakker and Daniel Tennant for their contributions to the Na_i metabolism studies.

Competing Interests

The Authors declare that there are no competing interests associated with the manuscript.

References

- 1 Shattock, M.J. (2009) Phospholemman: its role in normal cardiac physiology and potential as a drugable target in disease. *Curr. Opin. Pharmacol.* **9**, 160–166 <https://doi.org/10.1016/j.coph.2008.12.015>
- 2 Boguslavskiy, A., Pavlovic, D., Aughton, K., Clark, J.E., Howie, J., Fuller, W. et al. (2014) Cardiac hypertrophy in mice expressing unphosphorylatable phospholemman. *Cardiovasc. Res.* **104**, 72–82 <https://doi.org/10.1093/cvr/cvu182>

- 3 Okamoto, K., Wang, W., Rounds, J., Chambers, E.A. and Jacobs, D.O. (2001) ATP from glycolysis is required for normal sodium homeostasis in resting fast-twitch rodent skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* **281**, E479–E488 <https://doi.org/10.1152/ajpendo.2001.281.3.E479>
- 4 Cross, H.R., Radda, G.K. and Clarke, K. (1995) The role of Na⁺/K⁺ ATPase activity during low flow ischemia in preventing myocardial injury: a ³¹P, ²³Na and ⁸⁷Rb NMR spectroscopic study. *Magn. Reson. Med.* **34**, 673–685 <https://doi.org/10.1002/mrm.1910340505>
- 5 Murphy, E. and Eisner, D.A. (2009) Regulation of intracellular and mitochondrial sodium in health and disease. *Circ. Res.* **104**, 292–303 <https://doi.org/10.1161/CIRCRESAHA.108.189050>
- 6 Shattock, M.J. and Bers, D.M. (1989) Rat vs. rabbit ventricle: Ca flux and intracellular Na assessed by ion-selective microelectrodes. *Am. J. Physiol.* **256**, C813–C822 <https://doi.org/10.1152/ajpcell.1989.256.4.C813>
- 7 Kohlhaas, M., Liu, T., Knopp, A., Zeller, T., Ong, M.F., Bohm, M. et al. (2010) Elevated cytosolic Na⁺ increases mitochondrial formation of reactive oxygen species in failing cardiac myocytes. *Circulation* **121**, 1606–1613 <https://doi.org/10.1161/CIRCULATIONAHA.109.914911>
- 8 Liu, T. and O'Rourke, B. (2008) Enhancing mitochondrial Ca²⁺ uptake in myocytes from failing hearts restores energy supply and demand matching. *Circ. Res.* **103**, 279–288 <https://doi.org/10.1161/CIRCRESAHA.108.175919>
- 9 Harrison, S.M., McCall, E. and Boyett, M.R. (1992) The relationship between contraction and intracellular sodium in rat and guinea-pig ventricular myocytes. *J. Physiol.* **449**, 517–550 <https://doi.org/10.1113/jphysiol.1992.sp019100>
- 10 Despa, S., Islam, M.A., Weber, C.R., Pogwizd, S.M. and Bers, D.M. (2002) Intracellular Na⁺ concentration is elevated in heart failure but Na/K pump function is unchanged. *Circulation* **105**, 2543–2548 <https://doi.org/10.1161/01.CIR.0000016701.85760.97>
- 11 Neubauer, S., Newell, J.B. and Ingwall, J.S. (1992) Metabolic consequences and predictability of ventricular fibrillation in hypoxia. A ³¹P- and ²³Na-nuclear magnetic resonance study of the isolated rat heart. *Circulation* **86**, 302–310 <https://doi.org/10.1161/01.CIR.86.1.302>
- 12 Pogwizd, S.M., Sipido, K.R., Verdonck, F. and Bers, D.M. (2003) Intracellular Na in animal models of hypertrophy and heart failure: contractile function and arrhythmogenesis. *Cardiovasc. Res.* **57**, 887–896 [https://doi.org/10.1016/S0008-6363\(02\)00735-6](https://doi.org/10.1016/S0008-6363(02)00735-6)
- 13 Verdonck, F., Volders, P.G., Vos, M.A. and Sipido, K.R. (2003) Intracellular Na⁺ and altered Na⁺ transport mechanisms in cardiac hypertrophy and failure. *J. Mol. Cell. Cardiol.* **35**, 5–25 [https://doi.org/10.1016/S0022-2828\(02\)00280-8](https://doi.org/10.1016/S0022-2828(02)00280-8)
- 14 Fuller, W., Tulloch, L.B., Shattock, M.J., Calaghan, S.C., Howie, J. and Wypijewski, K.J. (2013) Regulation of the cardiac sodium pump. *Cell. Mol. Life Sci.* **70**, 1357–1380 <https://doi.org/10.1007/s00018-012-1134-y>
- 15 Pavlovic, D., Fuller, W. and Shattock, M.J. (2013) Novel regulation of cardiac Na pump via phospholemman. *J. Mol. Cell. Cardiol.* **61**, 83–93 <https://doi.org/10.1016/j.yjmcc.2013.05.002>
- 16 Shattock, M.J., Ottolia, M., Bers, D.M., Blaustein, M.P., Boguslavskyi, A., Bossuyt, J. et al. (2015) Na⁺/Ca²⁺ exchange and Na⁺/K⁺-ATPase in the heart. *J. Physiol.* **593**, 1361–1382 <https://doi.org/10.1113/jphysiol.2014.282319>
- 17 Pieske, B., Maier, L.S., Piacentino, Ill, V., Weisser, J., Hasenfuss, G. and Houser, S. (2002) Rate dependence of [Na⁺]_i and contractility in nonfailing and failing human myocardium. *Circulation* **106**, 447–453 <https://doi.org/10.1161/01.CIR.0000023042.50192.F4>
- 18 Verdonck, F., Volders, P.G., Vos, M.A. and Sipido, K.R. (2003) Increased Na⁺ concentration and altered Na/K pump activity in hypertrophied canine ventricular cells. *Cardiovasc. Res.* **57**, 1035–1043 [https://doi.org/10.1016/S0008-6363\(02\)00734-4](https://doi.org/10.1016/S0008-6363(02)00734-4)
- 19 Bossuyt, J., Ai, X., Moorman, J.R., Pogwizd, S.M. and Bers, D.M. (2005) Expression and phosphorylation of the Na-pump regulatory subunit phospholemman in heart failure. *Circ. Res.* **97**, 558–565 <https://doi.org/10.1161/01.RES.0000181172.27931.c3>
- 20 Murphy, E. and Allen, D.G. (2009) Why did the NHE inhibitor clinical trials fail? *J. Mol. Cell. Cardiol.* **46**, 137–141 <https://doi.org/10.1016/j.yjmcc.2008.09.715>
- 21 Balaban, R.S. (2002) Cardiac energy metabolism homeostasis: role of cytosolic calcium. *J. Mol. Cell. Cardiol.* **34**, 1259–1271 <https://doi.org/10.1006/jmcc.2002.2082>
- 22 Mootha, V.K., Arai, A.E. and Balaban, R.S. (1997) Maximum oxidative phosphorylation capacity of the mammalian heart. *Am. J. Physiol.* **272**, H769–H775 PMID:9124437
- 23 Neubauer, S. (2007) The failing heart — an engine out of fuel. *N. Engl. J. Med.* **356**, 1140–1151 <https://doi.org/10.1056/NEJMr063052>
- 24 Peterzan, M.A., Lygate, C.A., Neubauer, S. and Rider, O.J. (2017) Metabolic remodeling in hypertrophied and failing myocardium: a review. *Am. J. Physiol. Heart Circ. Physiol.* **313**, H597–H616 <https://doi.org/10.1152/ajpheart.00731.2016>
- 25 Gibbs, C.L. and Loiselle, D.S. (2001) Cardiac basal metabolism. *Jpn J. Physiol.* **51**, 399–426 <https://doi.org/10.2170/jphysiol.51.399>
- 26 Gibbs, C.L. (2003) Cardiac energetics: sense and nonsense. *Clin. Exp. Pharmacol. Physiol.* **30**, 598–603 <https://doi.org/10.1046/j.1440-1681.2003.03878.x>
- 27 Schramm, M., Klieber, H.G. and Daut, J. (1994) The energy expenditure of actomyosin-ATPase, Ca(2+)-ATPase and Na⁺,K(+) -ATPase in guinea-pig cardiac ventricular muscle. *J. Physiol.* **481**(Pt 3), 647–662 <https://doi.org/10.1113/jphysiol.1994.sp020471>
- 28 Stanley, W.C., Recchia, F.A. and Lopaschuk, G.D. (2005) Myocardial substrate metabolism in the normal and failing heart. *Physiol. Rev.* **85**, 1093–1129 <https://doi.org/10.1152/physrev.00006.2004>
- 29 Hue, L. and Taegtmeyer, H. (2009) The Randle cycle revisited: a new head for an old hat. *Am. J. Physiol. Endocrinol. Metab.* **297**, E578–E591 <https://doi.org/10.1152/ajpendo.00093.2009>
- 30 Aksentijević, D., McAndrew, D.J., Karlstädt, A., Zervou, S., Sebag-Montefiore, L., Cross, R. et al. (2014) Cardiac dysfunction and peri-weaning mortality in malonyl-coenzyme A decarboxylase (MCD) knockout mice as a consequence of restricting substrate plasticity. *J. Mol. Cell. Cardiol.* **75**, 76–87 <https://doi.org/10.1016/j.yjmcc.2014.07.008>
- 31 Kolwicz, S.C. and Tian, R. (2011) Glucose metabolism and cardiac hypertrophy. *Cardiovasc. Res.* **90**, 194–201 <https://doi.org/10.1093/cvr/cvr071>
- 32 Ventura-Clapier, R., Garnier, A. and Veksler, V. (2004) Energy metabolism in heart failure. *J. Physiol.* **555**, 1–13 <https://doi.org/10.1113/jphysiol.2003.055095>
- 33 Akki, A. and Seymour, A.-M.L. (2009) Western diet impairs metabolic remodelling and contractile efficiency in cardiac hypertrophy. *Cardiovasc. Res.* **81**, 610–617 <https://doi.org/10.1093/cvr/cvn316>
- 34 Chandler, M.P., Kerner, J., Huang, H., Vazquez, E., Reszko, A., Martini, W.Z. et al. (2004) Moderate severity heart failure does not involve a downregulation of myocardial fatty acid oxidation. *Am. J. Physiol. Heart Circ. Physiol.* **287**, H1538–H1543 <https://doi.org/10.1152/ajpheart.00281.2004>
- 35 Starling, R.C., Hammer, D.F. and Altschuld, R.A. (1998) Human myocardial ATP content and in vivo contractile function. *Mol. Cell. Biochem.* **180**, 171–177 <https://doi.org/10.1023/A:1006876031121>

- 36 Luptak, I., Sverdlov, A.L., Panagia, M., Qin, F., Pimentel, D.R., Croteau, D. et al. (2018) Decreased ATP production and myocardial contractile reserve in metabolic heart disease. *J. Mol. Cell. Cardiol.* **116**, 106–114 <https://doi.org/10.1016/j.yjmcc.2018.01.017>
- 37 Beer, M., Seyfarth, T., Sandstedt, J., Landschütz, W., Lipke, C., Köstler, H. et al. (2002) Absolute concentrations of high-energy phosphate metabolites in normal, hypertrophied, and failing human myocardium measured noninvasively with ^{31}P -SLOOP magnetic resonance spectroscopy. *J. Am. Coll. Cardiol.* **40**, 1267–1274 [https://doi.org/10.1016/S0735-1097\(02\)02160-5](https://doi.org/10.1016/S0735-1097(02)02160-5)
- 38 Garnier, A., Fortin, D., Deloménie, C., Momken, I., Veksler, V. and Ventura-Clapier, R. (2003) Depressed mitochondrial transcription factors and oxidative capacity in rat failing cardiac and skeletal muscles. *J. Physiol.* **551**, 491–501 <https://doi.org/10.1113/jphysiol.2003.045104>
- 39 Sharov, V.G., Todor, A.V., Silverman, N., Goldstein, S. and Sabbah, H.N. (2000) Abnormal mitochondrial respiration in failed human myocardium. *J. Mol. Cell. Cardiol.* **32**, 2361–2367 <https://doi.org/10.1006/jmcc.2000.1266>
- 40 Lemieux, H., Semsroth, S., Antretter, H., Höfer, D. and Gnaiger, E. (2011) Mitochondrial respiratory control and early defects of oxidative phosphorylation in the failing human heart. *Int. J. Biochem. Cell Biol.* **43**, 1729–1738 <https://doi.org/10.1016/j.biocel.2011.08.008>
- 41 Xu, K.Y., Zweier, J.L. and Becker, L.C. (1995) Functional coupling between glycolysis and sarcoplasmic reticulum Ca^{2+} transport. *Circ. Res.* **77**, 88–97 <https://doi.org/10.1161/01.RES.77.1.88>
- 42 Dizon, J., Burkhoff, D., Tauskela, J., Whang, J., Cannon, P. and Katz, J. (1998) Metabolic inhibition in the perfused rat heart: evidence for glycolytic requirement for normal sodium homeostasis. *Am. J. Physiol.* **274**, H1082–H1089 <https://doi.org/10.1152/ajpheart.1998.274.4.H1082>
- 43 Weiss, J. and Hittbrand, B. (1985) Functional compartmentation of glycolytic versus oxidative metabolism in isolated rabbit heart. *J. Clin. Invest.* **75**, 436–447 <https://doi.org/10.1172/JCI111718>
- 44 Fuller, W., Parmar, V., Eaton, P., Bell, J.R. and Shattock, M.J. (2003) Cardiac ischemia causes inhibition of the Na/K ATPase by a labile cytosolic compound whose production is linked to oxidant stress. *Cardiovasc. Res.* **57**, 1044–1051 [https://doi.org/10.1016/S0008-6363\(02\)00810-6](https://doi.org/10.1016/S0008-6363(02)00810-6)
- 45 Iwai, T., Tanonaka, K., Inoue, R., Kasahara, S., Motegi, K., Nagaya, S. et al. (2002) Sodium accumulation during ischemia induces mitochondrial damage in perfused rat hearts. *Cardiovasc. Res.* **55**, 141–149 [https://doi.org/10.1016/S0008-6363\(02\)00282-1](https://doi.org/10.1016/S0008-6363(02)00282-1)
- 46 Cox, D.A. and Matlib, M.A. (1993) Modulation of intramitochondrial free Ca^{2+} concentration by antagonists of Na^{+} - Ca^{2+} exchange. *Trends Pharmacol. Sci.* **14**, 408–413 [https://doi.org/10.1016/0165-6147\(93\)90063-P](https://doi.org/10.1016/0165-6147(93)90063-P)
- 47 Denton, R.M. and McCormack, J.G. (1990) Ca^{2+} as a second messenger within mitochondria of the heart and other tissues. *Annu. Rev. Physiol.* **52**, 451–466 <https://doi.org/10.1146/annurev.ph.52.030190.002315>
- 48 Eisner, D.A., Caldwell, J.L., Kistamas, K. and Trafford, A.W. (2017) Calcium and excitation–contraction coupling in the heart. *Circ. Res.* **121**, 181–195 <https://doi.org/10.1161/CIRCRESAHA.117.310230>
- 49 Hüser, J., Blatter, L.A. and Sheu, S.-S. (2000) Mitochondrial calcium in heart cells: beat-to-beat oscillations or slow integration of cytosolic transients? *J. Bioenerg. Biomembr.* **32**, 27–33 <https://doi.org/10.1023/A:1005556227425>
- 50 Maack, C., Cortassa, S., Aon, M.A., Ganesan, A.N., Liu, T. and O'Rourke, B. (2006) Elevated cytosolic Na^{+} decreases mitochondrial Ca^{2+} uptake during excitation-contraction coupling and impairs energetic adaptation in cardiac myocytes. *Circ. Res.* **99**, 172–182 <https://doi.org/10.1161/01.RES.0000232546.92777.05>
- 51 Eykyn, T.R., Aksentijević, D., Aughton, K.L., Southworth, R., Fuller, W. and Shattock, M.J. (2015) Multiple quantum filtered ^{23}Na NMR in the Langendorff perfused mouse heart: ratio of triple/double quantum filtered signals correlates with $[\text{Na}]_i$. *J. Mol. Cell. Cardiol.* **86**, 95–101 <https://doi.org/10.1016/j.yjmcc.2015.07.009>
- 52 Aksentijević, D., Karlstäedt, A., Basalay, M., O'Brien, B.A., Thakker, A., Tennant, D. et al. (2018) Causal link between intracellular Na overload and metabolic remodeling in the heart: uncoupling ATP supply and demand? *Heart* **104**, A2–A3 <https://doi.org/10.1136/heartjnl-2018-BSCR.6>
- 53 O'Brien, B.A., Aksentijević, D., Eykyn, T.R. and Shattock, M.J. (2015) Ouabain induces metabolic changes in the absence of contractile work in the blebbistatin-perfused rat heart. *J. Mol. Cell. Cardiol.* **86**, S47
- 54 Payne, G.S., Seymour, A.-M.L., Styles, P. and Radda, G.K. (1990) Multiple quantum filtered ^{23}Na NMR spectroscopy in the perfused heart. *NMR Biomed.* **3**, 139–146 <https://doi.org/10.1002/nbm.1940030307>
- 55 Dizon, J.M., Tauskela, J.S., Wise, D., Burkhoff, D., Cannon, P.J. and Katz, J. (1996) Evaluation of triple-quantum-filtered ^{23}Na NMR in monitoring of intracellular Na content in the perfused rat heart: comparison of intra- and extracellular transverse relaxation and spectral amplitudes. *Magn. Reson. Med.* **35**, 336–345 <https://doi.org/10.1002/mrm.1910350311>
- 56 Aksentijević, D., Eykyn, T.R., Fuller, W. and Shattock, M.J. (2015) Metabolic consequences of the acute and chronic myocardial sodium elevation. *J. Mol. Cell. Cardiol.* **86**, 1–78 <https://doi.org/10.1016/j.yjmcc.2015.07.025>
- 57 Mansor, L.S., Mehta, K., Aksentijević, D., Carr, C.A., Lund, T., Cole, M.A. et al. (2016) Increased oxidative metabolism following hypoxia in the type 2 diabetic heart, despite normal hypoxia signalling and metabolic adaptation. *J. Physiol.* **594**, 307–320 <https://doi.org/10.1113/JP271242>
- 58 Madelin, G., Lee, J.S., Regatte, R.R. and Jerschow, A. (2014) Sodium MRI: methods and applications. *Prog. Nucl. Magn. Reson. Spectrosc.* **79**, 14–47 <https://doi.org/10.1016/j.pnmrs.2014.02.001>
- 59 Kolmakova, E.V., Haller, S.T., Kennedy, D.J., Isachkina, A.N., Budny, G.V., Frolova, E.V. et al. (2011) Endogenous cardiotonic steroids in chronic renal failure. *Nephrol. Dial. Transplant.* **26**, 2912–2919 <https://doi.org/10.1093/ndt/gfq772>
- 60 Kennedy, D.J., Shrestha, K., Sheehey, B., Li, X.S., Guggilam, A., Wu, Y. et al. (2015) Elevated plasma marinobufagenin, an endogenous cardiotonic steroid, is associated with right ventricular dysfunction and nitrative stress in heart failure. *Circ. Heart Fail.* **8**, 1068–1076 <https://doi.org/10.1161/CIRCHEARTFAILURE.114.001976>
- 61 Gonano, L.A., Sepulveda, M., Rico, Y., Kaetzel, M., Valverde, C.A., Dedman, J. et al. (2011) Calcium-calmodulin kinase II mediates digitalis-induced arrhythmias. *Circ. Arrhythm. Electrophysiol.* **4**, 947–957 <https://doi.org/10.1161/CIRCEP.111.964908>
- 62 Matthews, P.M., Williams, S.R., Seymour, A.-M., Schwartz, A., Dube, G., Gadian, D.G. et al. (1982) A ^{31}P -NMR study of some metabolic and functional effects of the inotropic agents epinephrine and ouabain, and the ionophore R02-2985 (X537A) in the isolated, perfused rat heart. *Biochim. Biophys. Acta, Mol. Cell Res.* **720**, 163–171 [https://doi.org/10.1016/0167-4889\(82\)90008-8](https://doi.org/10.1016/0167-4889(82)90008-8)
- 63 Bersin, R.M., Wolfe, C., Kwassman, M., Lau, D., Klinski, C., Tanaka, K. et al. (1994) Improved hemodynamic function and mechanical efficiency in congestive heart failure with sodium dichloroacetate. *J. Am. Coll. Cardiol.* **23**, 1617–1624 [https://doi.org/10.1016/0735-1097\(94\)90665-3](https://doi.org/10.1016/0735-1097(94)90665-3)
- 64 Bersin, R.M. and Stacpoole, P.W. (1997) Dichloroacetate as metabolic therapy for myocardial ischemia and failure. *Am. Heart J.* **134**, 841–855 [https://doi.org/10.1016/S0002-8703\(97\)80007-5](https://doi.org/10.1016/S0002-8703(97)80007-5)

- 65 Fragasso, G., Pallosi, A., Puccetti, P., Silipigni, C., Rossodivita, A., Pala, M. et al. (2006) A randomized clinical trial of trimetazidine, a partial free fatty acid oxidation inhibitor, in patients with heart failure. *J. Am. Coll. Cardiol.* **48**, 992–998 <https://doi.org/10.1016/j.jacc.2006.03.060>
- 66 Tuunanen, H., Engblom, E., Naum, A., Nagren, K., Scheinin, M., Hesse, B. et al. (2008) Trimetazidine, a metabolic modulator, has cardiac and extracardiac benefits in idiopathic dilated cardiomyopathy. *Circulation* **118**, 1250–1258 <https://doi.org/10.1161/CIRCULATIONAHA.108.778019>
- 67 Gray, R.P., McIntyre, H., Sheridan, D.S. and Fry, C.H. (2001) Intracellular sodium and contractile function in hypertrophied human and guinea-pig myocardium. *Pflugers Arch.* **442**, 117–123 <https://doi.org/10.1007/s004240000512>
- 68 Eisner, D.A., Lederer, W.J. and Vaughan-Jones, R.D. (1984) The quantitative relationship between twitch tension and intracellular sodium activity in sheep cardiac Purkinje fibres. *J. Physiol.* **355**, 251–266 <https://doi.org/10.1113/jphysiol.1984.sp015417>
- 69 Lee, C.O., Kang, D.H., Sokol, J.H. and Lee, K.S. (1980) Relation between intracellular Na ion activity and tension of sheep cardiac Purkinje fibers exposed to dihydro-ouabain. *Biophys. J.* **29**, 315–330 [https://doi.org/10.1016/S0006-3495\(80\)85135-6](https://doi.org/10.1016/S0006-3495(80)85135-6)
- 70 Cohen, C.J., Fozzard, H.A. and Sheu, S.S. (1982) Increase in intracellular sodium ion activity during stimulation in mammalian cardiac muscle. *Circ. Res.* **50**, 651–662 <https://doi.org/10.1161/01.RES.50.5.651>
- 71 Lee, C.O. and Dagostino, M. (1982) Effect of strophanthidin on intracellular Na ion activity and twitch tension of constantly driven canine cardiac Purkinje fibers. *Biophys. J.* **40**, 185–198 [https://doi.org/10.1016/S0006-3495\(82\)84474-3](https://doi.org/10.1016/S0006-3495(82)84474-3)
- 72 Schmied, R., Wang, G.X. and Korth, M. (1991) Intracellular Na⁺ activity and positive inotropic effect of sulmazole in guinea pig ventricular myocardium. Comparison with a cardioactive steroid. *Circ. Res.* **68**, 597–604 <https://doi.org/10.1161/01.RES.68.2.597>
- 73 Wilde, A.A. and Kléber, A.G. (1986) The combined effects of hypoxia, high K⁺, and acidosis on the intracellular sodium activity and resting potential in guinea pig papillary muscle. *Circ. Res.* **58**, 249–256 <https://doi.org/10.1161/01.RES.58.2.249>
- 74 Wang, G.X., Schmied, R., Ebner, F. and Korth, M. (1993) Intracellular sodium activity and its regulation in guinea-pig atrial myocardium. *J. Physiol.* **465**, 73–84 <https://doi.org/10.1113/jphysiol.1993.sp019667>
- 75 Jelicks, L.A. and Siri, F.M. (1995) Effects of hypertrophy and heart failure on [Na⁺]_i in pressure-overloaded guinea pig heart. *Am. J. Hypertens.* **8**, 934–943 [https://doi.org/10.1016/0895-7061\(95\)00219-F](https://doi.org/10.1016/0895-7061(95)00219-F)
- 76 Liu, Y., Cabo, C., Salomonsz, R., Delmar, M., Davidenko, J. and Jalife, J. (1993) Effects of diacetyl monoxime on the electrical properties of sheep and guinea pig ventricular muscle. *Cardiovasc. Res.* **27**, 1991–1997 <https://doi.org/10.1093/cvr/27.11.1991>
- 77 Liu, T., Takimoto, E., Dimaano, V.L., DeMazumder, D., Kettlewell, S., Smith, G. et al. (2014) Inhibiting mitochondrial Na⁺/Ca²⁺ exchange prevents sudden death in a guinea pig model of heart failure. *Circ. Res.* **115**, 44–54 <https://doi.org/10.1161/CIRCRESAHA.115.303062>
- 78 Baudet, S., Noireaud, J. and Léoty, C. (1991) Intracellular Na activity measurements in the control and hypertrophied heart of the ferret: an ion-sensitive micro-electrode study. *Pflugers Arch.* **418**, 313–318 <https://doi.org/10.1007/BF00550867>
- 79 Levi, A.J., Lee, C.O. and Brooksby, P. (1994) Properties of the fluorescent sodium indicator 'SBFI' in rat and rabbit cardiac myocytes. *J. Cardiovasc. Electrophysiol.* **5**, 241–257 <https://doi.org/10.1111/j.1540-8167.1994.tb01161.x>
- 80 Grupp, I., Im, W.B., Lee, C.O., Lee, S.W., Pecker, M.S. and Schwartz, A. (1985) Relation of sodium pump inhibition to positive inotropy at low concentrations of ouabain in rat heart muscle. *J. Physiol.* **360**, 149–160 <https://doi.org/10.1113/jphysiol.1985.sp015609>
- 81 Donoso, P., Mill, J.G., O'Neill, S.C. and Eisner, D.A. (1992) Fluorescence measurements of cytoplasmic and mitochondrial sodium concentration in rat ventricular myocytes. *J. Physiol.* **448**, 493–509 <https://doi.org/10.1113/jphysiol.1992.sp019053>
- 82 Schepkin, V.D., Choy, I.O., Budinger, T.F., Obayashi, D.Y., Taylor, S.E., DeCampi, W.M. et al. (1998) Sodium TQF NMR and intracellular sodium in isolated crystalloid perfused rat heart. *Magn. Reson. Med.* **39**, 557–563 <https://doi.org/10.1002/mrm.1910390408>